

Numerical Solution of Nonlinear Singular Index-1 DAEs

A. Dick¹, O. Koch¹, R. März², and E. B. Weinmüller¹

Abstract: We discuss the convergence behavior of collocation schemes applied to approximate solutions of BVPs in nonlinear DAEs given in the form

$$f((D(t)x(t))', x(t), t) = 0, \quad t \in (0, 1], \quad B_0 D(0)x(0) + B_1 D(1)x(1) = \beta,$$

where $f(y, x, t) \in \mathbb{R}^m$, $D(t) \in \mathbb{R}^{n \times m}$, $y \in \mathbb{R}^n$, $x \in \mathcal{D}$, with $\mathcal{D} \subseteq \mathbb{R}^m$, $t \in [0, 1]$, $n \leq m$. Moreover, we require

$$\ker f_y(y, x, t) = 0, \quad (y, x, t) \in \mathbb{R}^n \times \mathcal{D} \times (0, 1], \quad \mathcal{R}(D(t)) = \mathbb{R}^n, \quad t \in [0, 1].$$

These conditions guarantee that the problem has a *properly stated leading term*. Note that at $t = 0$ the matrix $f_y(y, x, t)$ may undergo a rank drop. Let Q_0 be a continuous pointwise projector function onto $\ker D$ and let G_0 and G_1 be the matrices used in the decoupling of the DAE system into the inherent ODE system for Dx and the algebraic constraints, $(y, x, t) \in \mathbb{R}^n \times \mathcal{D} \times [0, 1]$,

$$G_0(y, x, t) := f_y(y, x, t)D(t), \quad G_1(y, x, t) := G_0(y, x, t) + f_x(y, x, t)Q_0(t).$$

We discuss DAE systems which are regular with tractability index 1 on $\mathbb{R}^n \times \mathcal{D} \times (0, 1]$. Consequently, $G_1(y, x, t)$ is nonsingular on $\mathbb{R}^n \times \mathcal{D} \times (0, 1]$. We permit a singular behavior of $G_1(y, x, t)$ for $t \rightarrow 0$, causing a singularity in the associated inherent ODE. To this end, we assume that $tG_1(y, x, t)^{-1}$ has a continuous extension on $\mathbb{R}^n \times \mathcal{D} \times [0, 1]$. The above assumptions result in the following form of the associated inherent ODE system, $u = Dx$,

$$u'(t) = \frac{1}{t}M(t)u(t) + h(u(t), t), \quad t \in (0, 1], \quad B_0 u(0) + B_1 u(1) = \beta, \quad u \in C[0, 1],$$

exhibiting a *singularity of the first kind*. We apply standard polynomial collocation at k interior collocation points to solve the *DAE system written in its original form*. We specify conditions under which for a certain class of well-posed boundary value problems in DAEs having sufficiently smooth solutions, the global error at the *collocation points* shows the convergence behavior $O(h^k)$ known from the ODE case. Here, the boundedness of the so-called canonical projector function plays a crucial role. The superconvergence order in the mesh points known for special choices of collocation points (Gaussian) for regular ODEs, does not hold even for singular ODEs, and thus neither for the DAEs, in general. The theoretical results are supported by numerical experiments.

¹ Institute for Analysis and Scientific Computing
Vienna University of Technology
Wiedner Hauptstrasse 8-10, A-1040 Wien, Austria
e.weinmueller@tuwien.ac.at, othmar@othmar-koch.org
e0425474@student.tuwien.ac.at

² Department of Mathematics Humboldt University Berlin
Rudower Chaussee 25, D-12489 Berlin, Germany
maerz@mathematik.hu-berlin.de